


# ELECTROGRAVITATION AND A NEW GRAVITATIONAL CONSTANT

BY

JERRY E. BAYLES



X,Y,Z

The following constants are pertinent to this paper and are all in the MKS system of units.

$m_e := 9.109389700 \cdot 10^{-31} \cdot \text{kg}$	Electron rest mass.
$q_o := 1.602177330 \cdot 10^{-19} \cdot \text{coul}$	Electron quantum charge.
$\mu_o := 1.256637061 \cdot 10^{-06} \cdot \text{henry} \cdot \text{m}^{-1}$	Magnetic permeability.
$\epsilon_o := 8.854187817 \cdot 10^{-12} \cdot \text{farad} \cdot \text{m}^{-1}$	Dielectric permittivity.
$r_c := 3.861593255 \cdot 10^{-13} \cdot \text{m}$	Compton electron radius.
$l_q := 2.817940920 \cdot 10^{-15} \cdot \text{m}$	Classic electron radius.
$c := 2.997924580 \cdot 10^{08} \cdot \text{m} \cdot \text{sec}^{-1}$	Speed of light in vacuum.
$\alpha := 7.297353080 \cdot 10^{-03}$	Fine structure constant.
$G := 6.672590000 \cdot 10^{-11} \cdot \text{newton} \cdot \text{m}^2 \cdot \text{kg}^{-2}$	Accepted gravitational constant.
$R_{n1} := 5.291772490 \cdot 10^{-11} \cdot \text{m}$	Bohr radius of Hydrogen.
$h := 6.626075500 \cdot 10^{-34} \cdot \text{joule} \cdot \text{sec}$	Plank constant.

These are the currently accepted values. The below constants are related directly to the theory of electrogravitation proposed by this author.

$V_{LM} := 8.542454612 \cdot 10^{-02} \cdot \text{m} \cdot \text{sec}^{-1}$	Least quantum velocity.
$f_{LM} := 1.003224805 \cdot 10^1 \cdot \text{Hz}$	Least quantum frequency.
$L_Q := 2.5729832158 \cdot 10^3 \cdot \text{henry}$	Least quantum inductance.
$C_Q := 3.861593281 \cdot 10^{-6} \cdot \text{farad}$	Least quantum capacitance.
$i_{LM} := q_o \cdot f_{LM}$ or, $i_{LM} = 1.60734404 \cdot 10^{-18} \cdot \text{amp}$ (= Least quantum amp.)	

The similar forms of the classical equations for the electric, magnetic, and gravitational forces suggest that they may be related to each other by a common mechanism of action. These are presented below as equations (1) through (3), respectively.

$$1) \quad F = \frac{q_0^2}{4 \cdot \pi \cdot \epsilon_0 \cdot r^2} \quad 2) \quad F = \frac{\mu_0 \cdot m_1 \cdot m_2}{4 \cdot \pi \cdot r^2} \quad 3) \quad F = \frac{G \cdot M_1 \cdot M_2}{r^2}$$

The terms  $m_1$  and  $m_2$  are the magnetic pole strengths in a classical magnetic force equation.  $M_1$  and  $M_2$  are the macroscopic mass terms in the classical gravitational force equation and  $q_0$  is the fundamental charge unit.

The purpose of this paper will be to establish a causal quantum mechanism for the gravitational force that will unify not only the electric and magnetic forces but the weak and strong forces also. This will be accomplished through the reduction of the electron field energy to its smallest quantum value. This value will then be utilized in a cross-product vector potential method to arrive at a solution for why the gravitational force is always one of attraction. The electron is considered in this paper to be a fundamental particle that has a direct connection to all of the known forces.

Furthermore, this paper will consider the existence of five fundamental forces instead of the normally assumed four forces. The electromagnetic, strong, weak, and gravitational forces will be expanded by considering that the electromagnetic force is composed of the electric and magnetic forces as individual action forces.

It is not the purpose of this paper to attempt to overthrow the concept of curved space as presented by Albert Einstein's General Theory Of Relativity but rather to propose that curved space is caused by the action of *electrogravitation*\*. Therefore, curved space is the result of gravitation and not the cause.

Recent experiments that attempted to refine the value of the accepted value of the gravitational constant have revealed a fairly large discrepancy, not only between the new values, but the old value as well. A quote from the April 29, 1995 issue of Science News is, "Now, experiments by three independent groups have produced values for the strength of the gravitational force (G) that disagree significantly with the currently accepted number and with each other." (See Reference [1], p. 15.)

Further, from the May 18, 1996 issue of Science News, "The news that three respected research groups had independently produced values for the strength of the gravitational force (G) that disagreed significantly with the currently accepted number and with each other created a considerable stir last year." (See Reference [2], p. 15.)

Finally, a quote from the March 1996 issue of Discover Magazine, "Ever since Isaac Newton watched an apple fall to the ground, scientists have taken gravity for granted. Until, that is, they tried to measure its strength with high-tech precision. Their results were so incredibly far off as to be newsworthy." (See Reference [3], p. 15.)

The results quoted above can be accounted for by the quantum vector potential nature of electrogravitation as proposed in this paper. It can be shown that the rest mass of the electron may be expressed by the charge couplet equation in (4) below. It is suggested that the geometry of the electron is made up of standing wave field energy caused by the uncertainty in the position of that electron (and its charge) which forms a right-handed triad in its quantum-jump actions.

$$4) \quad m_{\text{electron}} := \frac{\mu_0 \cdot q_0 \cdot q_0}{4 \cdot \pi \cdot l_q} \quad \text{or,} \quad m_{\text{electron}} = 9.10938969 \cdot 10^{-31} \cdot \text{kg}$$

This value is exact compared to the accepted rest mass of the electron, where:

$$m_e = 9.1093897 \cdot 10^{-31} \cdot \text{kg}$$

The quantum field energy that exists at the Compton radius of the electron is presented by equations (5) and (6) below. This electric field energy is equal to the rest mass of the electron times the fine structure constant.

$$5) \quad r_c := \frac{h}{2 \cdot \pi \cdot m_e \cdot c} \quad \text{or,} \quad r_c = 3.86159325 \cdot 10^{-13} \cdot \text{m}$$

$$6) \quad E_{fld} := \frac{q_o^2}{4 \cdot \pi \cdot \epsilon_o \cdot r_c} \quad \text{or,} \quad E_{fld} = 5.97442404 \cdot 10^{-16} \cdot \text{joule}$$

$$\text{Note that this is equal to:} \quad m_e \cdot c^2 \cdot \alpha = 5.97442409 \cdot 10^{-16} \cdot \text{joule}$$

The next equation (7) will define the least quantum mass related to the electric field energy of the electron presented by equation (6) above. This is the point where mass and energy are not just equivalent in magnitude, they are one and the same.

$$7) \quad m_{fld} := \frac{E_{fld}}{c^2} \quad \text{or,} \quad m_{fld} = 6.64744324 \cdot 10^{-33} \cdot \text{kg}$$

Then the least quantum field mass and energy are given in equation (8) as:

$$8) \quad e_{fld} := m_e \cdot \alpha \cdot \left( \frac{\text{m}}{\text{sec}} \right)^2 \quad \text{or,} \quad e_{fld} = 6.6474433 \cdot 10^{-33} \cdot \text{joule}$$

To say they are one and the same is to state that the least quantum mass and energy are interchangeable in an instant of time. A unit velocity implies the Dirac delta function at work which allows for the value of one as a limit concerning velocity times velocity. That is,  $(v1) \times (v2) = 1$  by  $\delta(v)$ .

A least quantum rotational velocity related directly to the electron rest mass may now be solved for. This is done in equation (9) below.

$$9) \quad V_{LM} := \sqrt{\frac{e_{fld}}{m_e}} \quad \text{or,} \quad V_{LM} = 0.08542455 \cdot \text{m} \cdot \text{sec}^{-1}$$

This is the equivalent linear circular field velocity in electron quantum space.

The next step will define the least quantum magnetic interaction energy. This is presented by equation (10) below.

$$10) \quad E_{\text{mag}} := m_e \cdot V_{\text{LM}}^2 \quad \text{or,} \quad E_{\text{mag}} = 6.6474433 \cdot 10^{-33} \cdot \text{joule}$$

It is now possible to present the electrogravitational least quantum energy equation below in equation (11) utilizing the result from equation (10) above.

$$11) \quad E_G := \left( \frac{\mu_o \cdot q_o^2}{4 \cdot \pi \cdot l_q} \cdot V_{\text{LM}}^2 \right) \cdot \mu_o \cdot \left( \frac{\mu_o \cdot q_o^2}{4 \cdot \pi \cdot l_q} \cdot V_{\text{LM}}^2 \right)$$

$$\text{Where; } E_G = 5.55289096 \cdot 10^{-71} \cdot \left( \frac{\text{henry}}{\text{m}} \right) \cdot \text{joule}^2$$

Separate system interaction at a distance is implied by equation (11) above.

The electrogravitational energy quantum above in equation (11) may be related to force. This is done in equation (12) below where the force between two electrons at the Bohr radius is found.

$$12) \quad F_{\text{Gnew}} := \frac{E_G}{R_{n1}^2} \quad \text{or,} \quad F_{\text{Gnew}} = 1.98297308 \cdot 10^{-50} \cdot \left( \frac{\text{henry}}{\text{m}} \right) \cdot \text{newton}^2$$

Compare this in magnitude and units to the classical value of gravitation obtained with equation (13) below.

$$13) \quad F_G := \frac{G \cdot m_e \cdot m_e}{R_{n1}^2} \quad \text{or,} \quad F_G = 1.97729139 \cdot 10^{-50} \cdot \text{newton}$$

The magnitudes are close but it is obvious that the units are not the same. It is suggested by this discrepancy that an important discovery may be found concerning the units not being identical since the magnitudes are so very close. In fact, a value for G may be found that incorporates these new units in equation (12) below.

Equation (10) previous may be employed to find the least quantum frequency.

This is presented by equation (14) below.

$$14) \quad f_{LM} := \frac{E_{\text{mag}}}{h} \quad \text{or,} \quad f_{LM} = 10.03224805 \cdot \text{Hz}$$

It may be further pointed out that several electrogravitational force equations can be presented that will all yield the same answers. This indicates that the gravitational force-field theory presented herein spans a great many of the forms of energy and force branches on the tree of physics. Three of those equations are presented below in equations (15), (16), and (17).

$$15) \quad F1_{Gnew} := \left( \frac{h \cdot f_{LM}}{R_{n1}} \right) \cdot \mu_o \cdot \left( \frac{h \cdot f_{LM}}{R_{n1}} \right)$$

$$\text{or,} \quad F1_{Gnew} = 1.98297308 \cdot 10^{-50} \cdot \frac{\text{henry}}{\text{m}} \cdot \text{newton}^2$$

$$16) \quad F2_{Gnew} := \left( \frac{L_{Q \cdot i_{LM}^2}}{R_{n1}} \right) \cdot \mu_o \cdot \left( \frac{L_{Q \cdot i_{LM}^2}}{R_{n1}} \right)$$

$$\text{or,} \quad F2_{Gnew} = 1.98297308 \cdot 10^{-50} \cdot \left( \frac{\text{henry}}{\text{m}} \right) \cdot \text{newton}^2$$

$$17) \quad F3_{Gnew} := \frac{m_e \cdot V_{LM}^2}{R_{n1}} \cdot \mu_o \cdot \frac{m_e \cdot V_{LM}^2}{R_{n1}}$$

$$\text{or,} \quad F3_{Gnew} = 1.98297308 \cdot 10^{-50} \cdot \left( \frac{\text{henry}}{\text{m}} \right) \cdot \text{newton}^2$$

It is easily seen that all three answers in equations (15), (16), and (17) are equal in magnitude and units.

It can also be shown that the famous Biot-Savart law that relates the magnetic field generated by a current can be incorporated into an electrogravitational expression also. This is presented by equation (20) on the following page.

First let us define the electrogravitational domain wavelength as:

$$18) \quad \lambda_{LM} := \frac{V_{LM}}{f_{LM}} \quad \text{or,} \quad \lambda_{LM} = 8.51499542 \cdot 10^{-3} \cdot \text{m}$$

Also let the following angles be defined:

$$19) \quad \theta := \frac{\pi}{2} \quad \text{and} \quad \phi := \frac{\pi}{2}$$

Then the Biot-Savart equation for the electrogravitational force between two electrons separated by the Bohr radius is given below in equation set (20).

$$20a) \quad F_{\text{sys1}} := \left( q_o \cdot V_{LM} \cdot \sin(\phi) \right) \cdot \left( \frac{\mu_o \cdot i_{LM} \cdot \lambda_{LM} \cdot \sin(\theta)}{4 \cdot \pi \cdot l_q \cdot R_{n1}} \right)$$

$$20b) \quad F_{\text{sys2}} := \left( q_o \cdot V_{LM} \cdot \sin(\phi) \right) \cdot \left( \frac{\mu_o \cdot i_{LM} \cdot \lambda_{LM} \cdot \sin(\theta)}{4 \cdot \pi \cdot l_q \cdot R_{n1}} \right)$$

Then finally;

20c)

$$F4_{G_{\text{new}}} := F_{\text{sys1}} \cdot \mu_o \cdot F_{\text{sys2}} \quad \text{or,} \quad F4_{G_{\text{new}}} = 1.98297308 \cdot 10^{-50} \cdot \frac{\text{henry}}{\text{m}} \cdot \text{newton}^2$$

The portion of the equations for the individual system forces that is the Biot-Savart least quantum expression at the Bohr radius is given below in equation (21).

$$21) \quad B_{LM} := \left( \frac{\mu_o \cdot i_{LM} \cdot \lambda_{LM} \cdot \sin(\theta)}{4 \cdot \pi \cdot l_q \cdot R_{n1}} \right) \quad \text{or,} \quad B_{LM} = 9.17825701 \cdot 10^{-3} \cdot \text{tesla}$$

Both of the equations in equation (20a,b) are of the standard form,  $F = qV \times \mathbf{B}$ .

Now we have enough of what may be called a preponderance of evidence that will support the case for assigning new units to the classic value of G. This new value is stated below in equation (22).

$$22) \quad G_{\text{new}} := \mu_o \cdot V_{LM}^4 \quad \text{or,} \quad G_{\text{new}} = 6.6917635 \cdot 10^{-11} \cdot \text{henry} \cdot \left( \frac{\text{m}^3}{\text{sec}^4} \right)$$

The ratio of this new proposed value of Gnew to G is:

$$23) \quad \frac{G_{\text{new}}}{G} = 1.00287347 \cdot \frac{\text{henry}}{\text{m}} \cdot \text{newton}$$

The new value of G may be inserted into the classical formula for the gravitational force and if we allow the expression for mass in equation (4) previous, then the result is an electrogravitational expression. This is presented in equation (24) below.

$$24) \quad F5_{Gnew} := \frac{(\mu_o \cdot V_{LM}^4) \cdot \left(\frac{\mu_o \cdot q_o^2}{4 \cdot \pi \cdot l_q}\right) \cdot \left(\frac{\mu_o \cdot q_o^2}{4 \cdot \pi \cdot l_q}\right)}{R_{n1}^2}$$

$$\text{or,} \quad F5_{Gnew} = 1.98297308 \cdot 10^{-50} \cdot \frac{\text{henry}}{\text{m}} \cdot \text{newton}^2$$

The above equation is now in the same general form as the classic gravitational expression. What is different are the extra henry/m and newton units. These 'extra' units are hidden units since the henry/m unit is a constant and the newton squared portion is actually inversely proportional to distance where one quantum newton force is also a constant. Thus on a macroscopic scale the simpler form of the classic gravitational force expression is assumed to be a correct form.

The following quote is from the book, Feynman Lectures on Gravitation, where Feynman's thoughts on the subject of the gravitational constant were condensed by the editor of the book, Brian Hatfield. He summed Feynman's conclusions as; "Of course, he expected that there might be difficulties in defining a consistent quantum theory (for example, the dimension of the gravitational constant is an obstacle to renormalization)." (See Reference [4], p. 15.)

It is suggested by this author that the problem of renormalization may be more easily solved by using the new value as defined in equation (22) previous.

The macroscopic electrogravitational case of the force of gravity between two massive objects may be obtained by increasing the magnitude of each force system by the ratio of each system mass to the rest mass of one electron. This is demonstrated using the energy value from equation (10) in equation (25) below.

First let the mass of the Earth and a mass on the surface of the Earth be defined as:

$$M_{\text{Earth}} := 5.98 \cdot 10^{24} \cdot \text{kg} \quad M_{\text{test}} := 1 \cdot \text{kg} \quad R_{\text{Earth}} := 6.37 \cdot 10^6 \cdot \text{m}$$

Then the ratios are:

$$R_1 := \frac{M_{\text{Earth}}}{m_e} \quad R_2 := \frac{M_{\text{test}}}{m_e}$$

Finally;

$$25) \quad F_{G_{\text{new}}} := \frac{R_1 \cdot E_{\text{mag}}}{R_{\text{Earth}}} \cdot \mu_0 \cdot \frac{R_2 \cdot E_{\text{mag}}}{R_{\text{Earth}}}$$

or,

$$F_{G_{\text{new}}} = 9.86195242 \cdot \frac{\text{henry}}{\text{m}} \cdot \text{newton}^2$$

Compare this with the classical gravitational expression in equation (26) below.

$$26) \quad F'_G := \frac{G \cdot M_{\text{Earth}} \cdot M_{\text{test}}}{R_{\text{Earth}}^2} \quad \text{or,} \quad F'_G = 9.83369558 \cdot \text{newton}$$

It is suggested by this author that the errors discovered in the recent attempts to measure the gravitational constant may be due to at least two effects. The first cause of error may be due to the metal and electronics that are part of the experimental hardware interacting with the quantum vector potentials generated in the Earth's molten core and stray ground currents associated with other actions near the Earth's surface. The second cause of error is that caused by the movement of mass in the locale of the test apparatus. It seems logical that if electrogravitation can cause a mass to accelerate, then accelerating a mass should create electrogravitation. (More specifically, a wave of gravitation.) This could be a stronger influence than that accounted for by ordinary gravitational influences since the electrogravitational wave would have a strength related to the rate of acceleration of the mass as well as the magnitude of the mass.

It is suggested by this author that sensitive quantum interference detectors feeding an amplifier tuned to  $f_{\text{LM}}$  might detect nearby mass accelerations.

One of the strongest arguments against an electromagnetic connection to the gravitational field was that an electromagnetic field can be shielded against while the gravitational field cannot. Further, the electromagnetic field has a bipolar aspect consisting of a negative and positive sense in the field and is a closed field such that all magnetic lines form a closed loop. The gravitational field apparently has no counterpart aspect of repulsion as does the magnetic or electric fields. The magnetic vector potential, (MVP), **can** however act through the best of shielding and when combined with the concept of the vector cross-product of two quantum uncertain currents acting 90 degrees to each other, the quantum electrogravitational action is generated that we take to be what is currently called gravity. Even though the action is unidirectional and always outwards from the origin, the reaction is a mirror image and is the conjugate of the action vector in every way. Thus, the total interaction that occurs partly in normal space is closed through the classic quantum radius points through imaginary energy space while to an outside observer in normal space it would appear that a monopole action had just occurred.


The Aharonov-Bohm effect has been demonstrated by actual experiment to prove that there exists quantum electromagnetic action through normally effective shielding.

The following is quoted from the April 1989 issue of Scientific American, (pages 56 to 62), "When the theories of relativity and quantum mechanics were introduced, the potentials, not the electric and magnetic fields, appeared in the equations of quantum mechanics, and the equations of relativity simplified into a compact mathematical form if the fields were expressed in terms of potentials." (See Reference [5], p. 15.). Also, "The consequence of the Aharonov-Bohm effect is that the potentials, not the fields, act directly on charges." (Reference [5], p. 15.)

The combination of the cross-product and the vector potential will serve to describe the electrogravitational action that will be presented on the next page.

First let the following drawing be offered to possibly help clarify the cross-product approach involving vector potentials that will soon be presented. This is labeled as figure #1 below.

Figure #1



System 1, rotation is Z into X

System 2, rotation is Z into -X

27)  $A1 := \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$      $B1 := \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$

28)  $A2 := \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$      $B2 := \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix}$

Vector cross-product of system 1 is:

Vector cross-product of system 2 is;

29)  $Sys1 := A1 \times B1$

30)  $Sys2 := A2 \times B2$

$$Sys1 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

$$Sys2 = \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix}$$

The total product of system 1 and 2 will yield the sign of the unit-scale electrogravitational action as:

31)  $F_{g1} := Sys1 \cdot Sys2$     or,     $F_{g1} = -1$

The above cross-product approach defines the action as a force of attraction by standard convention. The total interaction is independent of a preferred choice of axis or system since the reaction is always the conjugate of the action system. The result is always attraction between any interacting triad systems.

Another argument that is used as a reason for why the gravitational force cannot be attributed to any electrical force is that the mass of the proton is larger than the mass of the electron by a magnitude of 1836.152756 times that of the electron mass, while the charges are identical in magnitude. Therefore, if some kind of conversion constant is attempted utilizing only the charge values, the unequal mass values will introduce a very large error when the gravitational force is calculated using the classic gravitational equation.

It can be shown that not only can the above argument be rebuffed concerning the mass inequality versus charge equality but the relativistic aspect of mass may also be addressed as well.

First let the following constants be defined:

$$m_p := 1.672623100 \cdot 10^{-27} \cdot \text{kg} \quad = \text{proton rest mass.}$$

Then the least quantum radius associated with this mass is:

$$32) \quad l_{qp} := \frac{h \cdot \alpha}{2 \cdot \pi \cdot m_p \cdot c} \quad \text{or,} \quad l_{qp} = 1.53469853 \cdot 10^{-18} \cdot \text{m}$$

It is now pointed out that the relativistic aspect of mass may be stated below in equation (33).

$$33) \quad m'_p = \frac{m_p}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (\text{This will directly affect } l_{qp} \text{ above.})$$

The point of deriving the classical particle radius as a function of its relativistic mass is to indicate that the concepts presented by this author do not intend to divorce the theory as presented from the special or general laws of relativity but rather include Einstein's theory when relativistic velocities and large gravitational potential gradients are present.

The next page will further examine the electron-proton interaction in detail.

Let us establish the magnetic vectors for a proton-electron electrogravitational action at the  $r_{n1}$  radius of the Bohr atom of Hydrogen. First system 1 is established as:

$$34) \quad \text{icp1}_a := \frac{q_o \cdot V_{LM}}{I_{qp}} \cdot \sin(\theta) \quad \text{and,} \quad \text{icp1}_b := \frac{q_o \cdot V_{LM}}{R_{n1}} \cdot \sin(\phi)$$

where;

$$\text{icp1}_a = 8.91805577 \cdot 10^{-3} \cdot \text{amp} \quad \text{and,} \quad \text{icp1}_b = 2.5863786 \cdot 10^{-10} \cdot \text{amp}$$

Then the magnetic vectors associated with the above current potentials are;

$$35) \quad \text{Acp1} := \begin{pmatrix} 0 \\ 0 \\ \text{icp1}_a \end{pmatrix} \cdot \text{amp} \quad \text{Bcp1} := \begin{pmatrix} \text{icp1}_b \\ 0 \\ 0 \end{pmatrix} \quad \text{Rotation is:} \\ \text{Z into X.}$$

Then inserting the correct dimensional constants into the cross-product of the current potentials above;

$$36) \quad \text{Sys1}_{cp} := \frac{\mu_o}{4 \cdot \pi} \cdot (\text{Acp1} \times \text{Bcp1}) \quad \text{This is the proton triad system} \\ \text{outgoing vector potential.}$$

$$\text{or,} \quad \text{Sys1}_{cp} = \begin{pmatrix} 0 \\ 2.30654686 \cdot 10^{-19} \\ 0 \end{pmatrix} \cdot \text{newton}$$

The electron triad vector potential system is now calculated beginning with the statement for the A & B vectors which shall be labeled as Sys2.

$$37) \quad \text{icp2}_a := \frac{q_o \cdot V_{LM}}{I_q} \cdot \sin(\theta) \quad \text{and,} \quad \text{icp2}_b := \frac{q_o \cdot V_{LM}}{R_{n1}} \cdot \sin(\phi)$$

where;

$$\text{icp2}_a = 4.85692479 \cdot 10^{-6} \cdot \text{amp} \quad \text{and} \quad \text{icp2}_b = 2.5863786 \cdot 10^{-10} \cdot \text{amp}$$

Then the magnetic vectors associated with the above current potentials are:

$$38) \quad \text{Acp2} := \begin{pmatrix} \text{icp2}_a \\ 0 \\ 0 \end{pmatrix} \quad \text{Bcp2} := \begin{pmatrix} 0 \\ 0 \\ \text{icp2}_b \end{pmatrix} \cdot \text{amp} \quad \text{Rotation is:} \\ \text{X into Z.}$$

Then again inserting the correct dimensional constants for the electron triad cross-product of the current potentials in equation (38) previous;

$$39) \quad \text{Sys2}_{cp} := \frac{\mu_o}{4 \cdot \pi} \cdot (\text{Acp2} \times \text{Bcp2}) \quad \text{This is the electron triad system outgoing vector potential.}$$

$$\text{or,} \quad \text{Sys2}_{cp} = \begin{pmatrix} 0 \\ -1.25618463 \cdot 10^{-22} \\ 0 \end{pmatrix} \cdot \text{newton}$$

Then the total electrogravitational interaction force between a proton and an electron at the  $r_{n1}$  orbital of the element Hydrogen is;

$$40) \quad F_{gep} := \text{Sys1}_{cp} \cdot \mu_o \cdot \text{Sys2}_{cp} \quad \text{or,} \quad F_{gep} = -3.64104145 \cdot 10^{-47} \cdot \text{newton}^2 \cdot \left( \frac{\text{henry}}{\text{m}} \right)$$

Let us now calculate the classic gravitational force for the same parameters involving a proton and electron mass separated by the Bohr radius:

$$41) \quad F_G := \frac{G \cdot m_p \cdot m_e}{R_{n1}^2} \quad \text{or,} \quad F_G = 3.63060903 \cdot 10^{-47} \cdot \text{newton}$$

It is to be noted that charge polarity is not a factor since a (+) charge going in a given direction has the B field given as conforming to the right-hand rule and thus the force vector potential is in the same direction as a (-) charge going in the same direction as the (+) charge but has the B field going in a direction opposite to the right-hand rule. Thus the charge polarity is arbitrary and only the fact that vector potential forces are based on the right-hand triad system as previously presented need be considered in their calculation.

The above counters one of the common arguments against electromagnetic forces being applicable to the gravitational action due to the fact that the electron-proton force is different than the electron-electron force at the same considered distance.

The weak and strong forces may be shown below in equations (42) and (43). They follow the same general form as the electrogravitational equation.

$$42) \quad F_w = \frac{q_o^2}{4 \cdot \pi \cdot \epsilon_o \cdot R_x^2} \cdot \left( \frac{\pi}{\epsilon_o} \right) \cdot \left( \frac{\mu_o \cdot q_o^2 \cdot V \cdot LM^2}{4 \cdot \pi \cdot r_c \cdot R_x} \right) = \text{weak force equation.}$$

$$43) \quad F_s = \frac{q_o^2}{4 \cdot \pi \cdot \epsilon_o \cdot R_x^2} \cdot \left( \frac{2 \cdot \pi \cdot R_{n1}}{\epsilon_o \cdot R_x} \right) \cdot \left( \frac{\mu_o \cdot q_o^2 \cdot V \cdot LM^2}{4 \cdot \pi \cdot r_c \cdot R_x} \right) = \text{strong force equation.}$$

If the proper nuclear  $R_x$  (radius) values are plugged into equations (42) and (43) above the resulting forces will be close to the presently accepted values. They also may be calculated using the cross-product vector potential approach as outlined in this paper in previous equations.

In concluding this paper this author would like to say that while the classical gravitational equation was the first equation to be formalized concerning force at a distance, it has stubbornly refused to be improved upon with the possible exception of Einstein's General Theory of Relativity. Unfortunately this theory has not explained the mechanics correctly or we would have solved the anti-gravity puzzle. This paper is a new approach utilizing the very basic accepted classical equations as a starting point to put the gravitational action in a logical engineering format and at the same time in terms of the more recent formulas of quantum physics. -- Jerry E. Bayles

END

REFERENCES

1. Ivars Peterson, "Gravity's force: Chasing an elusive constant," *Science News*, 29 April 1995: 263.
2. Ivars Peterson, "Measuring the gravitational constant," *Science News*, 18 May 1996: 319.
3. Hans Christian von Baeyer, "Big G," *Discover*, March 1996: 96.
4. Richard P. Feynman, Fernando B. Moringo, and William G. Wagner, *Feynman Lectures on Gravitation* (Menlo Park, Calif.: Addison-Wesley Publishing Company, June 1995), xxxi.
5. Yoseph Imry and Richard A. Webb, "Quantum Interference and the Aharonov-Bohm Effect," *Scientific American*, April 1989: 56.